

Strange axial-vector mesons mixing angle

De-Min Li*, Zhen Li

Department of Physics, Zhengzhou University, Zhengzhou, Henan 450052, P. R. China

February 2, 2008

Abstract

The masses of the $K_1(^3P_1)$ and $K_1(^1P_1)$ are considered in a nonrelativistic constituent quark model, and the absolute value of the $K_1(^3P_1) - K_1(^1P_1)$ mixing angle is determined to be about 59.29° . Comparison of the theoretical predictions on the strong decay widths of the $K_1(1270)$ and $K_1(1400)$ in the 3P_0 decay model as well as the production ratio of these two states in the τ decay between the available experimental data strongly favors that the $K_1(^3P_1) - K_1(^1P_1)$ mixing angle is about $+59.29^\circ$.

Key words: Other strange mesons; Nonrelativistic quark model; Hadronic decays of mesons; Decays of taus

PACS numbers:14.40.Ev; 12.39.Jh; 13.25.-k; 13.35.Dx

*E-mail: lidm@zzu.edu.cn

1 Introduction

The strange axial vector mesons provide interesting possibilities to study the QCD in the nonperturbative regime by the mixing of the 3P_1 and 1P_1 states. In the exact SU(3) limit, the $K_1(^3P_1)$ and $K_1(^1P_1)$ do not mix, just as the a_1 and b_1 mesons do not mix. For the strange quark mass greater than the up and down quark masses so that SU(3) is broken, also, the $K_1(^3P_1)$ and $K_1(^1P_1)$ do not possess definite C-parity, therefore these states can in principle mix to give the physical $K_1(1270)$ and $K_1(1400)$.

Accurate determination of θ_K , the mixing angle of the $K_1(^3P_1)$ and $K_1(^1P_1)$, is important for comparing the theory predictions about the decays involving the strange axial-mesons with the experimental data. In the literature, θ_K has been estimated by some different approaches, however, there is not yet a consensus on the value of θ_K . As the optimum fit to the data as of 1977, Carnegie et al. finds $\theta_K = (41 \pm 4)^\circ$ [1]. Within the heavy quark effective theory Isgur and Wise predict two possible mixing angles, $\theta_K \sim 35.3^\circ$ and $\theta_K \sim -54.7^\circ$ [2]. Based on the analysis of $\tau \rightarrow \nu K_1(1270)$ and $\tau \rightarrow \nu K_1(1400)$, Rosner suggests $\theta_K \sim 62^\circ$ [3], Asner et al. gives $\theta_K = (69 \pm 16 \pm 19)^\circ$ or $(49 \pm 16 \pm 19)^\circ$ [4], and Cheng obtains $\theta_K = \pm 37^\circ$ or $\pm 58^\circ$ [5]. From the experimental information on masses and the partial rates of $K_1(1270)$ and $K_1(1400)$, Suzuki finds two possible solutions with a two-fold ambiguity, $\theta_K \sim 33^\circ$ or 57° [6]. A constraint $35^\circ \leq \theta_K \leq 55^\circ$ is predicted by Burakovsky et al. in a nonrelativistic constituent quark model[7], and within the same model, the values of $\theta_K \simeq (31 \pm 4)^\circ$ and $\theta_K \simeq (37.3 \pm 3.2)^\circ$ are suggested by Chliapnikov[8] and Burakovsky[9], respectively. The calculations for the strong decays of $K_1(1270)$ and $K_1(1400)$ in the 3P_0 decay model suggest $\theta_K \sim 45^\circ$ [10, 11]. The mixing angles $\theta_K \sim 34^\circ$ [12], $\theta_K \sim 5^\circ$ [13] are also presented within a relativized quark model. Vijande et al. suggests $\theta_K \sim 55.7^\circ$ based on the calculations in a constituent quark model[14]. More recently, based on the $f_1(1285) - f_1(1420)$ mixing angle $\sim 50^\circ$ derived from the analysis for a substantial body of data concerning the $f_1(1420)$ and $f_1(1285)$ [15], we suggest that the $K_1(^3P_1) - K_1(^1P_1)$ mixing angle is about $\pm(59.55 \pm 2.81)^\circ$ [16].

In the present work, we shall show that the $K_1(^3P_1) - K_1(^1P_1)$ mixing angle derived from the nonrelativistic constituent quark model is in good agreement with that given by Ref.[16], and

try to constrain the sign of the $K_1(^3P_1) - K_1(^1P_1)$ mixing angle by considering the open-flavor strong decays of the $K_1(1270)$ and $K_1(1400)$ in the 3P_0 decay model and the production ratio of these two states in the τ decay.

2 Nonrelativistic constituent quark model for P -wave mesons

In the constituent quark model, the conventional $q\bar{q}$ wave function is typically assumed to be a solution of a nonrelativistic Schrödinger equation with the generalized Breit-Fermi Hamiltonian which contains a QCD inspired potential $V(\mathbf{r})$ [17]. The phenomenological forms of the matrix element of the Breit-Fermi Hamiltonian for the $q\bar{q}$ mesons with orbital angular momentum L are given by[8, 18]:

$$M_{L=0} = m_q + m_{\bar{q}} + e_0 \frac{\langle \mathbf{s}_q \cdot \mathbf{s}_{\bar{q}} \rangle}{m_q m_{\bar{q}}}, \quad (1)$$

$$\begin{aligned} M_{L \neq 0} = & m_q + m_{\bar{q}} + a_L + b_L \left(\frac{1}{m_q} + \frac{1}{m_{\bar{q}}} \right) + c_L \left(\frac{1}{m_q^2} + \frac{1}{m_{\bar{q}}^2} \right) + \frac{d_L}{m_q m_{\bar{q}}} + e_L \frac{\langle \mathbf{s}_q \cdot \mathbf{s}_{\bar{q}} \rangle}{m_q m_{\bar{q}}} \\ & + f_L \left(\frac{1}{m_q^3} + \frac{1}{m_{\bar{q}}^3} \right) + g_L \left[\frac{(m_q + m_{\bar{q}})^2 + 2m_q m_{\bar{q}}}{4m_q^2 m_{\bar{q}}^2} \langle \mathbf{L} \cdot \mathbf{S} \rangle - \frac{m_q^2 - m_{\bar{q}}^2}{4m_q^2 m_{\bar{q}}^2} \langle \mathbf{L} \cdot \mathbf{S}_- \rangle \right] \\ & + \frac{h_L}{m_q m_{\bar{q}}} \langle \mathbf{S}_{q\bar{q}} \rangle, \end{aligned} \quad (2)$$

where m_q and $m_{\bar{q}}$ are the constituent quark masses, $\mathbf{s}_q$ and $\mathbf{s}_{\bar{q}}$ are the constituent quark spins, e_0 , a_L , b_L , c_L , d_L , e_L , f_L , g_L and h_L are constants, $\mathbf{S} = \mathbf{s}_q + \mathbf{s}_{\bar{q}}$, $\mathbf{S}_- = \mathbf{s}_q - \mathbf{s}_{\bar{q}}$, and $\mathbf{S}_{q\bar{q}} = 3 \frac{(\mathbf{s}_q \cdot \mathbf{r})(\mathbf{s}_{\bar{q}} \cdot \mathbf{r})}{r^2} - \mathbf{s}_q \cdot \mathbf{s}_{\bar{q}}$. Angular momentum part of the matrix elements of (1) and (2) is shown in Table 1.

	3P_2	3P_1	3P_0	1P_1	3S_1	1S_0
$\langle \mathbf{s}_q \cdot \mathbf{s}_{\bar{q}} \rangle$	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{4}$	$-\frac{3}{4}$	$\frac{1}{4}$	$-\frac{3}{4}$
$\langle \mathbf{L} \cdot \mathbf{S} \rangle$	1	-1	-2	0		
$\langle \mathbf{S}_{q\bar{q}} \rangle$	$-\frac{2}{5}$	2	-4	0		
$\langle \mathbf{L} \cdot \mathbf{S}_- \rangle$	0	0	0	$\frac{3}{2}$		

Table 1: Angular momentum part of the matrix elements of (1) and (2).

With the help of Table 1, applying (1) and (2) to S -wave and P -wave mesons, in the SU(2)

flavor symmetry limit, one can obtain¹

$$\frac{M_\pi + 3M_\rho}{2M_K + 6M_{K^*} - M_\pi - 3M_\rho} = \frac{m_u}{m_s} = 0.6298 \pm 0.00068, \quad (3)$$

and

$$\frac{M(^3P_2)_{s\bar{s}} - M(^1P_1)_{s\bar{s}}}{M(^3P_2)_{n\bar{n}} - M(^1P_1)_{n\bar{n}}} = \frac{m_u^2}{m_s^2}. \quad (4)$$

From (4), with the help of the Gell-Mann-Okubo mass formula[20]

$$M^2(^3P_2)_{s\bar{s}} + M^2(^3P_2)_{n\bar{n}} = 2M_{K(^3P_2)}^2, \quad (5)$$

$$M^2(^1P_1)_{s\bar{s}} + M^2(^1P_1)_{n\bar{n}} = 2M_{K_1(^1P_1)}^2, \quad (6)$$

taking $M(^3P_2)_{n\bar{n}} = M_{a_2(1320)} = 1318.3 \pm 0.6$ MeV, $M(^1P_1)_{n\bar{n}} = M_{b_1(1235)} = 1229.5 \pm 3.2$ MeV and $M_{K(^3P_2)} = M_{K_2^*(1430)} = 1429 \pm 0.99$ MeV, one can arrive that

$$M_{K(^1P_1)} = 1369.52 \pm 1.92 \text{ MeV}. \quad (7)$$

The $K_1(^3P_1)$ and $K_1(^1P_1)$ can mix to produce the physical states $K_1(1400)$ and $K_1(1270)$ and the mixing between $K_1(^3P_1)$ and $K_1(^1P_1)$ can be parameterized as[6]

$$\begin{aligned} K_1(1400) &= K_1(^3P_1) \cos \theta_K - K_1(^1P_1) \sin \theta_K, \\ K_1(1270) &= K_1(^3P_1) \sin \theta_K + K_1(^1P_1) \cos \theta_K, \end{aligned} \quad (8)$$

where θ_K denotes the $K_1(^3P_1) - K_1(^1P_1)$ mixing angle. Without any assumption about the origin of the $K_1(^3P_1) - K_1(^1P_1)$ mixing, the masses of the $K_1(^3P_1)$ and $K_1(^1P_1)$ can be related to $M_{K_1(1400)}$ and $M_{K_1(1270)}$, the masses of the $K_1(1400)$ and $K_1(1270)$, by the following relation phenomenologically,

$$S \begin{pmatrix} M_{K_1(^3P_1)}^2 & A \\ A & M_{K_1(^1P_1)}^2 \end{pmatrix} S^\dagger = \begin{pmatrix} M_{K_1(1400)}^2 & 0 \\ 0 & M_{K_1(1270)}^2 \end{pmatrix}, \quad (9)$$

where A denotes a parameter describing the $K_1(^3P_1) - K_1(^1P_1)$ mixing, and

$$S = \begin{pmatrix} \cos \theta_K & -\sin \theta_K \\ \sin \theta_K & \cos \theta_K \end{pmatrix}.$$

¹Where $n\bar{n} = (u\bar{u} + d\bar{d})/\sqrt{2}$. All the masses used as input in the present work are taken from PDG[19].

From (9), one can have

$$M_{K_1(^3P_1)}^2 = M_{K_1(1400)}^2 \cos^2 \theta_K + M_{K_1(1270)}^2 \sin^2 \theta_K, \quad (10)$$

$$M_{K_1(^1P_1)}^2 = M_{K_1(1400)}^2 \sin^2 \theta_K + M_{K_1(1270)}^2 \cos^2 \theta_K, \quad (11)$$

$$\cos(2\theta_K) = \frac{M_{K_1(^3P_1)}^2 - M_{K_1(^1P_1)}^2}{M_{K_1(1400)}^2 - M_{K_1(1270)}^2}. \quad (12)$$

Inputting $M_{K_1(1400)} = 1402 \pm 7$ MeV, $M_{K_1(1270)} = 1273 \pm 7$ MeV and $M_{K_1(^3P_1)} \simeq 1369.52 \pm 1.92$ MeV shown in (7), from (10)-(12), we have

$$M_{K_1(^3P_1)} = 1307.88 \pm 10.33 \text{ MeV}, \quad \theta_K = \pm(59.29 \pm 2.87)^\circ. \quad (13)$$

Obviously, the present result that $(M_{K_1(^1P_1)}, M_{K_1(^3P_1)}) = (1369.5 \pm 1.92, 1307.88 \pm 10.33)$ MeV and $\theta_K = \pm(59.29 \pm 2.87)^\circ$ is in good agreement with that $(M_{K_1(^1P_1)}, M_{K_1(^3P_1)}) = (1370.03 \pm 9.69, 1307.35 \pm 0.63)$ MeV and $\theta_K = \pm(59.55 \pm 2.81)^\circ$ given by Ref.[16] based on the $f_1(1285) - f_1(1420)$ mixing angle $\sim 50^\circ$ extracted from the analysis for a substantial body of data concerning the $f_1(1420)$ and $f_1(1285)$ [15].

Within the nonrelativistic constituent quark model, the results regarding the masses of the $K_1(^1P_1)$ and $K_1(^3P_1)$, $(M_{K_1(^1P_1)}, M_{K_1(^3P_1)}) = (1368, 1306)$ MeV suggested by [8] and $(M_{K_1(^1P_1)}, M_{K_1(^3P_1)}) = (1356, 1322)$ MeV suggested by [9], are in good agreement with our predicted result. However, based on the following relation employed by [8, 9]

$$\tan^2(2\theta_K) = \left(\frac{M_{K_1(^3P_1)}^2 - M_{K_1(^1P_1)}^2}{M_{K_1(1400)}^2 - M_{K_1(1270)}^2} \right)^2 - 1, \quad (14)$$

the values of $\theta_K = (31 \pm 4)^\circ$ given by[8] and $\theta_K = (37.3 \pm 3.2)^\circ$ given by[9] disagree with value of $|\theta_K| \simeq (59.29 \pm 2.87)^\circ$ given by the present work.

As pointed out by our previous paper [16], (14) is equivalent to (12), and will yield two solutions $|\theta_K|$ and $\frac{\pi}{2} - |\theta_K|$. Simultaneously considering the relations (10), (11) and (14), in the presence of $M_{K_1(1400)} > M_{K_1(1270)}$, we can conclude that if $M_{K_1(^3P_1)} < M_{K_1(^1P_1)}$, the $|\theta_K|$ would greater than 45° . In fact, relation (12) clearly indicates that in the presence of $M_{K_1(1400)} > M_{K_1(1270)}$, the case $M_{K_1(^3P_1)} < M_{K_1(^1P_1)}$ must require $45^\circ < |\theta_K| < 90^\circ$.

3 The sign of θ_K constrained by experimental information

Now we wish to discuss the sign of θ_K by considering the open-flavor strong decays of the

$K_1(1270)$ and $K_1(1400)$ in the 3P_0 decay model, and the production ratio of these two physical strange states in the τ decay.

3.1 Strong decays of the $K_1(1270)$ and $K_1(1400)$ in the 3P_0 model

The main assumption of the 3P_0 decay model is that the strong decays take place via the production of a quark-antiquark pair with the vacuum quantum numbers which corresponds to the 3P_0 state of a quark-antiquark pair. After the 3P_0 decay model was originally introduced by Micu[21], it was applied extensively to meson and baryon decays. It is widely accepted that the 3P_0 model is successful since it gives a good description of many of the observed decay amplitudes and partial widths of the open-flavor meson strong decays.

Assuming a fixed 3P_0 source strength, simple harmonic oscillator quark model meson wave functions and physical phase space, Ackleh et al.[22] developed a diagrammatic, momentum-space formulation of the 3P_0 model to evaluate the partial width $\Gamma_{A \rightarrow BC}$

$$\Gamma_{A \rightarrow BC} = 2\pi \frac{PE_BE_C}{M_A} \sum_{LS} |\mathcal{M}_{LS}|^2, \quad (15)$$

where P is the decay momentum for the decay $A \rightarrow B + C$, E_B and E_C are the energies of mesons B and C , in the rest frame of A ,

$$\begin{aligned} P &= \frac{[(M_A^2 - (M_B + M_C)^2)(M_A^2 - (M_B - M_C)^2)]^{1/2}}{2M_A}, \\ E_B &= (M_A^2 - M_C^2 + M_B^2)/2M_A, \\ E_C &= (M_A^2 - M_B^2 + M_C^2)/2M_A, \end{aligned}$$

M_A , M_B and M_C denote the masses of the mesons A , B and C , respectively; $\mathcal{M}_{LS}$ are proportional to an overall Gaussian in $x = P/\beta$ times a channel-dependent polynomial $\mathcal{P}_{LS}$,

$$\mathcal{M}_{LS} = \frac{\gamma}{\pi^{1/4} \beta^{1/2}} \mathcal{P}_{LS}(x) e^{-x^2/12}.$$

It is found that this formulation with the width parameter $\beta = 0.4$ GeV and the pair-production strength parameter $\gamma = 0.4$ can give a reasonably accurate description of the overall scale of decay widths[23, 24].

Based on (8) and (15), employing the analytical results for $\mathcal{P}_{LS}$ listed in Appendix A of Ref.[23], one can have[24]

$$\Gamma(K_1(1270) \rightarrow \rho K) = 21.8 \cos^2 \theta_K + 61.6 \sin \theta_K \cos \theta_K + 43.6 \sin^2 \theta_K, \quad (16)$$

$$\Gamma(K_1(1270) \rightarrow \pi K^*) = 59.6 \cos^2 \theta_K - 158.7 \sin \theta_K \cos \theta_K + 115.7 \sin^2 \theta_K, \quad (17)$$

$$\Gamma_{\text{thy}}(K_1(1270)) = 81 \cos^2 \theta_K - 97 \sin \theta_K \cos \theta_K + 159 \sin^2 \theta_K, \quad (18)$$

$$\Gamma(K_1(1400) \rightarrow \rho K) = 160 \cos^2 \theta_K - 219.9 \sin \theta_K \cos \theta_K + 82.3 \sin^2 \theta_K, \quad (19)$$

$$\Gamma(K_1(1400) \rightarrow \omega K) = 52.3 \cos^2 \theta_K - 72.3 \sin \theta_K \cos \theta_K + 26.8 \sin^2 \theta_K, \quad (20)$$

$$\Gamma(K_1(1400) \rightarrow \pi K^*) = 141.1 \cos^2 \theta_K + 176.2 \sin \theta_K \cos \theta_K + 78.8 \sin^2 \theta_K, \quad (21)$$

$$\Gamma_{\text{thy}}(K_1(1400)) = 353 \cos^2 \theta_K - 116 \sin \theta_K \cos \theta_K + 188 \sin^2 \theta_K, \quad (22)$$

$$|D/S|^2 = \begin{cases} \frac{(-0.0411 \cos \theta_K - 0.029 \sin \theta_K)^2}{(-0.204 \cos \theta_K + 0.288 \sin \theta_K)^2}, & \text{for } K_1(1270) \rightarrow \pi K^* \\ \frac{(-0.0498 \cos \theta_K + 0.0704 \sin \theta_K)^2}{(+0.247 \cos \theta_K + 0.175 \sin \theta_K)^2}, & \text{for } K_1(1400) \rightarrow \pi K^* \end{cases}. \quad (23)$$

For $\theta_K = \pm(59.29 \pm 2.87)^\circ$, the theoretical results regarding the above widths are shown in Tables 2, 3 and 4. Tables 2-4 clearly indicate that the present experimental data strongly prefer $\theta_K = +(59.29 \pm 2.87)^\circ$ over $\theta_K = -(59.29 \pm 2.87)^\circ$.

$K_1(1270)$	Exp.[19]	$\theta_K = +(59.29 \pm 2.87)^\circ$	$\theta_K = -(59.29 \pm 2.87)^\circ$
Γ (MeV)	90±20	96.07 ± 5.76	181.25 ± 1.11
$\Gamma(\rho K)/\Gamma(\pi K^*)$	2.625±0.902	2.07 ± 0.41	0.064 ± 0.014

Table 2: The predicted results of the $K_1(1270)$ strong decays in the 3P_0 decay model.

$K_1(1400)$	Exp.[19]	$\theta_K = +(59.29 \pm 2.87)^\circ$	$\theta_K = -(59.29 \pm 2.87)^\circ$
Γ (MeV)	174±13	180.1 ± 4.48	282.0 ± 10.0
$\Gamma(\rho K)/\Gamma$	0.03±0.03	0.033 ± 0.01	0.71 ± 0.04
$\Gamma(\omega K)/\Gamma$	0.01±0.01	0.0095 ± 0.0034	0.23 ± 0.01
$\Gamma(\pi K^*)/\Gamma$	0.94±0.06	0.96 ± 0.05	0.063 ± 0.006

Table 3: The predicted results of the $K_1(1400)$ strong decays in the 3P_0 decay model.

$ D/S ^2$	Exp.[19]	$\theta_K = +(59.29 \pm 2.87)^\circ$	$\theta_K = -(59.29 \pm 2.87)^\circ$
$K_1(1270) \rightarrow \pi K^*$	1.0$\pm$0.7	0.1 ± 0.03	0.0001 ± 0.0002
$K_1(1400) \rightarrow \pi K^*$	0.04$\pm$0.01	0.02 ± 0.004	12.5 ± 15.6

Table 4: The $|D/S|^2$ ratios for $K_1(1270) \rightarrow \pi K^*$ and $K_1(1400) \rightarrow \pi K^*$ in the 3P_0 model.

3.2 Production ratio of the $K_1(1270)$ and $K_1(1400)$ in the τ decay

With the definition of the decay constant of the axial-vector meson given by[5]

$$\langle 0 | A_\mu | A(q, \varepsilon) \rangle = f_A m_A \varepsilon_\mu, \quad (24)$$

the partial width for $\tau \rightarrow \nu_\tau K_1$ can be expressed by

$$\Gamma(\tau \rightarrow \nu_\tau K_1) = \frac{G_F^2}{16\pi} |V_{us}|^2 f_{K_1}^2 \frac{(m_\tau^2 + 2m_{K_1}^2)(m_\tau^2 - m_{K_1}^2)^2}{m_\tau^3}. \quad (25)$$

Considering the SU(3) breaking corrections, following Ref.[5, 6], we have

$$\frac{f_{K_1(1270)} m_{K_1(1270)}}{f_{K_1(1400)} m_{K_1(1400)}} = \frac{\sin \theta_K - \delta \cos \theta_K}{\cos \theta_K + \delta \sin \theta_K}, \quad (26)$$

where the parameter δ denoting a SU(3) breaking factor has the following form in the static limit of the quark model[10]²

$$\delta = \frac{1}{\sqrt{2}} \frac{m_s - m_u}{m_s + m_u} = 0.16 \pm 0.0003. \quad (27)$$

From (25) and (26), the $K_1(1400)$ and $K_1(1270)$ production ratio in the τ decay can be given by

$$\frac{\Gamma(\tau \rightarrow \nu_\tau K_1(1270))}{\Gamma(\tau \rightarrow \nu_\tau K_1(1400))} = F_p \left| \frac{\sin \theta_K - \delta \cos \theta_K}{\cos \theta_K + \delta \sin \theta_K} \right|^2, \quad (28)$$

where F_p denotes the phase factor given by

$$F_p = \frac{\left(m_\tau^2 + 2m_{K_1(1270)}^2 \right) \left(m_\tau^2 - m_{K_1(1270)}^2 \right)^2 m_{K_1(1400)}^2}{\left(m_\tau^2 + 2m_{K_1(1400)}^2 \right) \left(m_\tau^2 - m_{K_1(1400)}^2 \right)^2 m_{K_1(1270)}^2} = 1.82 \pm 0.086.$$

Then, from (28), one can have

$$\frac{\Gamma(\tau \rightarrow \nu_\tau K_1(1270))}{\Gamma(\tau \rightarrow \nu_\tau K_1(1400))} = \begin{cases} 2.62 \pm 0.55, & \text{for } \theta_K = +(59.29 \pm 2.87)^\circ \\ 11.59 \pm 3.43, & \text{for } \theta_K = -(59.29 \pm 2.87)^\circ \end{cases}. \quad (29)$$

² $m_u = 307.8 \pm 0.19$ MeV and $m_s = 488.69 \pm 0.28$ MeV derived from (1).

Experimentally, the $\mathcal{B}(\tau \rightarrow \nu_\tau K_1(1270))$ and $\mathcal{B}(\tau \rightarrow \nu_\tau K_1(1400))$ have been reported by TPC/Two-Gamma collaboration[25] in 1994 and ALEPH collaboration[26] in 1999, respectively. The averaged result of these two collaborations is given by[19]

$$\begin{aligned}\mathcal{B}(\tau \rightarrow \nu_\tau K_1(1270)) &= (0.47 \pm 0.11) \times 10^{-2} \\ \mathcal{B}(\tau \rightarrow \nu_\tau K_1(1400)) &= (0.17 \pm 0.26) \times 10^{-2}\end{aligned}\quad , \quad (30)$$

which gives that

$$\left. \frac{\Gamma(\tau \rightarrow \nu_\tau K_1(1270))}{\Gamma(\tau \rightarrow \nu_\tau K_1(1400))} \right|_{\text{exp-1}} = 2.76 \pm 4.28. \quad (31)$$

This measured result also is in favor of $\theta_K = +(59.29 \pm 2.87)^\circ$ over $\theta_K = -(59.29 \pm 2.87)^\circ$, although the uncertainty of the reported result is large as shown in (31).

Assuming the resonance structure of $\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$ decays being dominated by the $K_1(1270)$ and $K_1(1400)$ resonances, in 2000, both CLEO collaboration[4] and OPAL collaboration[27] have also measured the ratio of $\nu_\tau K_1(1270)$ to $\nu_\tau K_1(1400)$ with the averaged result[19]

$$\frac{\Gamma(\tau \rightarrow \nu_\tau K_1(1270))}{\Gamma(\tau \rightarrow \nu_\tau K_1(1270)) + \Gamma(\tau \rightarrow \nu_\tau K_1(1400))} = 0.69 \pm 0.15, \quad (32)$$

which therefore in turn implies that

$$\left. \frac{\Gamma(\tau \rightarrow \nu_\tau K_1(1270))}{\Gamma(\tau \rightarrow \nu_\tau K_1(1400))} \right|_{\text{exp-2}} = 2.23 \pm 1.56. \quad (33)$$

Comparison of (29) and (33) again shows that the present experimental data strongly prefer $\theta_K = +(59.29 \pm 2.87)^\circ$ over $\theta_K = -(59.29 \pm 2.87)^\circ$.

It is based on the $K_1(1400)$ production dominance in the τ decay that Suzuki suggests that the preferred result is $\theta_K \approx 33^\circ$ rather than 57° [6]. However, the recent available experiment data shown in (33) clearly show the $K_1(1270)$ dominance in the τ decay. Consequently, the argument of ruling out $\theta_K \approx 57^\circ$ from the $K_1(1400)$ dominance is therefore no longer valid. The study of hadronic decays $D \rightarrow K_1(1270)\pi, K_1(1400)\pi$ decays performed by Cheng[5] favors $\theta_K \approx -58^\circ$, however as pointed out by Cheng et al. in Ref.[28] that this argument is subject to many uncertainties such as the unknown $D \rightarrow K_1(^1P_1), K_1(^3P_1)$ transition form factors and the decay constants of $K_1(1270)$ and $K_1(1400)$. We note that the recent analysis for the SU(3) nonets of the axial vector mesons into a vector and a pseudoscalar performed by Roca et al.[29] based on a tensor formulation of the vector and axial vector fields gives $\theta_K = +(62 \pm 3)^\circ$, which is in fact in good agreement with our suggested result that $\theta_K = +(59.29 \pm 2.87)^\circ$.

4 Concluding remarks

In the nonrelativistic constituent quark model, the masses of the $K_1(^3P_1)$ and $K_1(^1P_1)$ are determined to be 1307.88 ± 10.33 and 1396.5 ± 1.92 MeV, respectively, which therefore suggests that the absolute value of the $K_1(^3P_1) - K_1(^1P_1)$ mixing angle is $(59.29 \pm 2.87)^\circ$. These findings are in good agreement with those given by Ref.[16] based on the investigation on the implication of the $f_1(1285) - f_1(1420)$ mixing for the $K_1(^3P_1) - K_1(^1P_1)$ mixing angle. Investigating the open-flavor strong decays of the $K_1(1270)$ and $K_1(1400)$ in the 3P_0 decay model, we find the current experimental data strongly prefer $\theta_K = +(59.29 \pm 2.87)^\circ$ over $\theta_K = -(59.29 \pm 2.87)^\circ$. The analysis for the production ratio of the $K_1(1270)$ and $K_1(1400)$ in the τ decay also indicates that the experimental data is in favor of the result $\theta_K = +(59.29 \pm 2.87)^\circ$.

In the framework of a covariant light-front quark model, the calculations performed by Cheng et al.[28] for the exclusive radiative B decays, $B \rightarrow K_1(1270)\gamma$, $K_1(1400)\gamma$, show that the relative strength of $B \rightarrow K_1(1270)\gamma$ and $B \rightarrow K_1(1400)\gamma$ rates is very sensitive to the sign of the θ_K . The recent analysis of two-body B decays with an axial-vector meson in the final state performed by Nardulli et al.[30, 31] using naive factorization, shows the branching ratios for $B \rightarrow b_1\pi$, b_1K , $a_1\pi$ and a_1K also depend strongly on the θ_K . In addition, as pointed by Suzuki[32], the relation $|Am(J/\psi(\psi') \rightarrow K_1^0(1400)\overline{K}^0)|^2 = \tan^2 \theta_K |Am(J/\psi(\psi') \rightarrow K_1^0(1270)\overline{K}^0)|^2$ can be able to determine the θ_K directly without referring to other parameters. Therefore, in order to further check the consistency of our suggested mixing angle of $K_1(1270)$ and $K_1(1400)$, detailed experimental study of the above mentioned decays involving the axial-vector mesons is certainly desirable.

Acknowledgments: This work is supported in part by National Natural Science Foundation of China under Contract No. 10205012, Henan Provincial Science Foundation for Outstanding Young Scholar under Contract No. 0412000300, Henan Provincial Natural Science Foundation under Contract No. 0311010800, Foundation of the Education Department of Henan Province under Contract No. 2003140025, and the Program for Youthful Excellent Teachers in University of Henan Province.

References

- [1] R. K. Carnegie et al, Phys. Lett. B **68**, 289 (1977)
- [2] N. Isgur, M. B. Wise, Phys. Lett. B **232**, 113 (1989)
- [3] J. Rosner, Com. Nucl. Part. Phys. **16**, 109 (1986)
- [4] D. M. Anser et al., CLEO collaboration, Phys. Rev. D **62**, 072006 (2000)
- [5] H. Y. Cheng, Phys. Rev. D **67**, 094007 (2003)
- [6] M. Suzuki, Phys. Rev. D **47**, 1252 (1997)
- [7] L. Burkovsky, T. Goldman, Phys. Rev. D **56**, 1368 (1997)
- [8] P. V. Chliapnikov, Phys. Lett. B **496**, 129 (2000)
- [9] L. Burakovsky, T. Goldman, Phys. Rev. D **57**, 2879 (1998)
- [10] H. G. Blundell, S. Godfrey, B. Phelps, Phys. Rev. D **53**, 3712 (1996)
- [11] T. Barnes, N. Black, P. R. Page, Phys. Rev. D **68**, 054014 (2003)
- [12] S. Godfrey, N. Isgur, Phys. Rev. D **32**, 189 (1985)
- [13] S. Godfrey, R. Kokoski, Phys. Rev. D **43**, 1679 (1991)
- [14] J. Vijande, F. Fernandez, A. Valcarce, J. Phys. G **31**, 481 (2005)
- [15] F. E. Close, A. Kirk, Z. Phys. C **76**, 469 (1997)
- [16] D. M. Li, B. Ma, H. Yu, Eur. Phys. J. A **26**, 141 (2005)
- [17] W. Lucha, F. F. Schöberl, D. Gromes, Phys. Rep. **200**, 127 (1991)
- [18] D. Flamm, F. Schöberl, Introduction to quark model of elementary particles, Vol. 1, Gordon and Breach Science Publishers, London, 1982
- [19] S. Eidelman et al., Phys. Lett. B **592**, 1 (2004)
- [20] S. Okubo, Prog. Theor. Phys. **27**, 949 (1962)
- [21] L. Micu, Nucl. Phys. B **10**, 521 (1969)
- [22] E. S. Ackleh, T. Barnes, Phys. Rev. D **54**, 6811 (1996)
- [23] T. Barnes, F. E. Close, P. R. Page, E. S. Swanson, Phys. Rev. D **55**, 4157 (1997)
- [24] T. Barnes, N. Black, P. R. Page, Phys. Rev. D **68**, 054014 (2003)

- [25] D. A. Bauer et al., TPC/Two-Gamma collaboration, Phys. Rev. D **50**, R13 (1994)
- [26] R. Barate et al., ALEPH collaboration, Eur. Phys. J. C **11**, 599 (1999)
- [27] G. Abbiendi et al., OPAL collaboration, Eur. Phys. J. C **13**, 197 (2000)
- [28] H. Y. Cheng, C. K. Chua, Phys. Rev. D **69**, 094007 (2004)
- [29] L. Roca, J. E. Palomar, E. Oset, Phys. Rev. D **70**, 094006 (2004)
- [30] G. Nardulli, T. N. Pham, Phys. Lett. B **623**, 65 (2005)
- [31] V. Laporta, G. Nardulli, T. N. Pham, hep-ph/0602243
- [32] M. Suzuki, Phys. Rev. D **55**, 2840 (1997)